

CHILD HEALTH CLINIC



Children with high risk or problems will require more services

SERVICE	2 MOS	4 MOS	6 MOS	9 MOS	12 MOS	18 MOS	2 YRS	3 YRS	4 YRS	5 YRS	6 YRS	9,12,15& 18 YRS
Health Hx.	X	X	X	X	X	X	X	X	X	X	X	X
Temperature	X	X	X	X	X	X	X	X	X	X	X	X
Length/Height	X	X	X	X	X	X	X	X	X	X	X	X
Weight	X	X	X	X	X	X	X	X	X	X	X	X
Head Circ.	X	X	X	X	X	X	X					
Blood Pressure								X	X	X	X	X
Physical Exam	X	X	X	X	X	X	X	X	X	X	X	X
Oral Screening	X	X	X	X	X	X	X	X	X	X	X	X
Dentist Referral	UPON ERUPTION OF FIRST TOOTH AND ROUTINELY AS NEEDED											
Dental Varnish	RECOMMENDED EVERY 3 TO 6 MONTHS AFTER ERUPTION OF FIRST TOOTH UP TO AGE 3 1/2 YRS (41 MOS)											
Vision								X	X	X	X	X
Hearing			X		X		X	X	X	X	X	X
Develop Screen			X		X		X	X	X	X		X
Nutritional Assess	X	X	X	X	X	X	X	X	X	X	X	X
Hemoglobin										X		
Urinalysis	INDICATED CLINICALLY											
Lead Screening					X		X					
Sickle Cell	WITH PARENTAL CONSENT AS INDICATED											
Anticipatory Guid.	X	X	X	X	X	X	X	X	X	X	X	X
Referral/Follow-Up	X	X	X	X	X	X	X	X	X	X	X	X

